

NYMphaeETUM CANDIDAE MILJAN 1933 IN POLAND

KRZYSZTOF SPAŁEK

Laboratory of Geobotany and Plant Conservation
Department of Biosystematics, University of Opole
Oleska 22, 45-052 Opole, Poland
e-mail: kspalek@uni.opole.pl

(Received: February 12, 2008. Accepted: June 12, 2008)

ABSTRACT

This paper presents the distribution, floristic composition and ecological requirements of the water plant community *Nymphaeetum candidae* Miljan 1933, belonging to the Potametea class. Until present this community has been reported only from north-eastern Poland. During present studies it was stated also in the south-western part of the country. *Nymphaeetum candidae* is considered to be a rare and endangered plant community in several countries of Central Europe.

KEY WORDS: *Nymphaeetum candidae*, Potametea class, distribution, phytosociology, endangered association, Poland.

INTRODUCTION

Biodiversity in Poland, measured at the level of species diversity, is among the highest in Central Europe. Also at the level of diversity of plant communities and wildlife habitats it is exceptionally rich, with e.g. well preserved wetlands (Andrzejewski and Weigle 1993). At present natural lakes, and in areas where they are lacking – ponds and oxbow lakes, are the most valuable in terms of preservation of rare and endangered plant communities (Wojtaszek 1989; Szumiec 1995; Spałek 2004, 2005, 2006a, b, 2007). One of the rarest aquatic plant communities in Poland is *Nymphaeetum candidae* Miljan 1933 (syn.: *Potamogetono-Nymphaeetum candidae* Hejný 1948 nom. invalid., *Nymphaeetum albo-candidae* (Hejný 1950) Passarge 1957), belonging to the Potametea class. Phytocoenoses of this community were first described from Estonia (Miljan 1933). *Nymphaeetum candidae* occurs in various types of shallow (50-150 cm) dystrophic, oligotrophic and mesotrophic (pH 5.0-6.0), stagnating or slowly running waters. Most often it inhabits lakes, oxbow lakes, ponds, channels and river bends of slow streams. This community is noted in many countries of Central Europe (e.g. Miljan 1933; Hejný 1948; Passarge 1957, 1996; Neuhäusl and Neuhäuslova 1965; Weber-Oldecop 1975; Hejný and Husák 1978; Jeckel 1981; Cernohous and Husak 1986; Pott 1995; Schubert et al. 1995; Dierssen 1996; Matuszkiewicz 2005; Balevičienė and Balevičius 2006). The characteristic and dominant species of *Nymphaeetum candidae* is *Nymphaea candida*. This association is considered to be poor in species and consists

of about 1-10 taxa on the average (Miljan 1933; Passarge 1957; Neuhäusl and Neuhäuslova 1965; Hejný and Husák 1978). Species such as *Potamogeton natans*, *Nuphar lutea*, *Batrachium circinatum*, *Juncus bulbosus* and *Utricularia minor* attain a higher constancy index (Miljan 1933; Passarge 1957; 1996; Neuhäusl and Neuhäuslova 1965; Weber-Oldecop 1975; Hejný and Husák 1978; Jeckel 1981; Cernohous and Husak 1986; Pott 1995). Until present only two sub-communities of this association: *Nymphaeetum candidae typicum* and *Nymphaeetum candidae elodeetosum* with the occurrence of *Elodea canadensis* and *Potamogeton lucens* (Passarge 1996) have been distinguished.

Nymphaea candida is an Euro-Siberian element of water flora with the western limit of the range on the line of the Rhine (Meusel et al. 1965; Hulten and Fries 1986; Muntendam et al. 1996; Wayda 2000). There is still a great deal of confusion considering its southern limit. Wayda (2000) suggested that *Nymphaea candida* occurs only in north-eastern Poland (Zajac and Zajac 2001). However, this species has also been reported from Czech Republic (Neuhäusl and Tomsovic 1957; Hejný and Slavik 1997) and from southern regions of Germany (Benkert et al. 1996). In Slovakia *Nymphaea candida* have not been recorded so far. According to Muntendam et al. (1996) and Tutin et al. (2002) *Nymphaea candida* also occurs in eastern France, Switzerland, south-western Romania, Austria, Hungary and former Yugoslavia. This species has been reported several times from the area of Silesia in south-western Poland (Fiek 1881; Schube 1903; Zabawski and Matuła 1973; Kłosow-

ski 2001). Unfortunately, there is no herbarium documentation from stations of this region. Recently there have been conducted studies aimed at identification of distribution of *Nymphaea candida* in some regions of south-western Poland (Nowak and Nowak 2007). Some of the examined specimens had all characters (Muntendam et al. 1996; Kubát 2002) of *Nymphaea candida*.

In Poland phytocoenoses of *Nymphaeetum candidae* are known mainly from lakes of clean, soft and mesotrophic, rarer – eutrophic waters, oxbow lakes and artificial water reservoirs in northern and eastern part of Poland (Tomaszewicz 1977, 1979; Mamiński 1986; Boiński 1993; Kłosowski and Tomaszewicz 1993; Sugier and Popiótek 1995, 1998). Exceptionally, locations of this community were reported from oxbow lakes of the Oder in Lower Silesia in south-western Poland (Macicka-Pawlik and Wilczyńska 1996). *Nymphaeetum candidae* occurs in water bodies of mineral and organic substratum. An increase of proportion of calcareous gyttja in sediments causes its retreat from inhabited localities (Kłosowski and Tomaszewicz 1993). In the phytosociological literature the community with domination of *Nymphaea candida* is described also as a variety or a facies within *Myriophyllo-Nupharetum* (Fijałkowski 1959; Podbielkowski 1968; Krzywański 1974). It was described as an association with *Nymphaea candida* by Hereźniak (1972).

This aquatic plant community is recognized as endangered at both regional and Central European scale (Hejny 1990; Kłosowski and Tomaszewicz 1993, Pott 1995; Schubert et al. 1995; Rennwald 2000).

This paper describes *Nymphaeetum candidae* community. The phytosociological table is given and floristic composition and ecological requirements of this community are discussed.

METHODS

The fieldwork was conducted during vegetation seasons 2000 and 2007. *Nymphaeetum candidae* community was studied following the Zurich-Montpellier School of Phytosociology (Braun-Blanquet 1964). Phytosociological nomenclature and syntaxonomical attachment are based on Oberdorfer (1994), Pott (1995) and Matuszkiewicz (2005). Vascular plant species names are given according to Mirek et al. (2002), and bryophytes to Frahm and Frey (1992). Hydrogen ion concentration was measured with Elmetron pH microcomputer CP-315.

RESULTS

During geobotanical investigations carried out in south-western Poland ten new localities of *Nymphaeetum candidae* were discovered (Fig. 1). Phytocoenoses of this community were formed usually in oxbow lakes of the Odra river – Stare Kolnie, Skorogoszcz, Stobrawa, Wielopole, ponds – Przysiecz, Sowin, Stary Węgliniec, Staw Nowokuznicki Nature Reserve, Winna Góra and exceptionally in small ponds in forest on the peatbog substratum – Zielonka. Localities in Stary Węgliniec and Zielonka are located in the vicinity of the continuous range of the species, covering south-eastern Germany (Meusel et al. 1965; Benkert et al. 1996). At newly discovered localities in south-western Poland *Nymphaeetum candidae* developed in transparent, eutrophic or mesotrophic waters, on sand or sand and silt basis, of depth 30-150 cm and pH 6.3-7.5 (Table 1). Phytocoenoses of this community covered areas from ca 50 m² up to 1 ha. In most patches this community has one-layer, rarer two-layers structure. In the water surface layer *Nymphaea candida* predominates, with coverage at the level



Fig. 1. Localities of *Nymphaeetum candidae* in Poland. ● – new locality, ○ – locality from the literature.

TABLE 1. The physico-chemical characteristics of water.

Locality	Cl [mg/l]	Electrolytic conductivity [$\mu\text{S}/\text{cm}^2$]	pH H ₂ O	BOD ₅ [mgO ₂ /l]	O ₂ Dissolved oxygen [mg/l]	PO ₄ Phosphates [mg/l]	Nitrate nitrogen [mg/l]	Ammonium nitrogen [mg/l]	Nitrite nitrogen [mg/l]
Przysiecz	8.32	89.76	7.1	0.91	4.38	0.11	0.35	0.57	0.014
Skorogoszcz	14.8	118.33	7.2	0.71	6.91	0.02	0.37	0.02	0.002
Sowin	8.23	90.01	7.2	0.98	5.45	0.19	0.42	1.35	0.016
Stare Kolnie	9.12	94.76	7.2	0.84	4.87	0.04	0.52	0.59	0.011
Staw Nowokuźnicki	9.27	89.97	7.1	0.89	6.68	0.13	0.34	1.03	0.016
Stary Węgliniec	9.11	117.24	7.2	0.70	6.98	0.02	0.32	0.03	0.003
Stobrawa	8.12	89.8	7.2	0.94	4.37	0.11	0.37	2.37	0.016
Wielopole	8.45	96.32	7.3	1.04	4.57	0.12	0.41	1.39	0.023
Winna Góra	9.19	96.95	6.9	0.95	4.78	0.10	0.44	0.58	0.017
Zielonka	10.45	96.80	6.3	2.02	5.29	0.12	0.47	0.55	0.013

35-70% (Table 2). In patches from 3 to 11, on average – 6 taxa were recorded. In total, in its phytocoenoses 15 plant species were noted. These patches should be recognised as belonging to the sub-community *Nymphaetum candidae* typicum (Passarge 1996). The floristic composition of patches of *Nymphaetum candidae* in south-western Poland is usually very similar to phytocoenoses of this community from other areas of Poland (Mamiński 1986; Boiński 1993; Kłosowski and Tomaszewicz 1993; Sugier and Popiołek 1995, 1998) and neighbouring countries (Miljan 1933; Passarge 1957, 1996; Neuhäusl and Neuhäuslova 1965; Weber-Oldecop 1975; Hejny and Husák 1978; Jeckel 1981;

Cernohous and Husak 1986). *Nymphaetum candidae* usually built a micro-mosaic complex with associations from classes: Potametea (e.g. *Potametum natantis*) and *Utricularietea intermedio-minoris* (e.g. *Sparganietum minimi*). In the processes of natural succession phytocoenoses of *Nymphaetum candidae* most often undergo transformation into *Scirpetum lacustris*, *Typhetum latifoliae*, *Caricetum rostratae*, *Equisetetum fluviatilis* and other rush associations. In the newly found localities in Poland phytocoenoses of *Nymphaetum candidae* occurred in a close contact with *Caricetum rostratae*, *Equisetetum fluviatilis* and *Phragmitetum australis*. Naturally, in the case of intensive fish far-

TABLE 2. *Nymphaetum candidae* Miljan 1933.

Relevé number	1	2	3	4	5	6	7	8	9	10	C
Date: year	2000	2002	2002	2003	2004	2005	2005	2006	2007	2007	
month	07	08	08	07	08	07	07	08	07	07	
day	04	11	11	23	09	30	18	17	16	18	
Locality	SN	WG	St	So	W	P	SW	Z	S	SK	
Cover of herb layer [%]	65	55	50	60	50	55	50	65	50	70	
Cover of moss layer [%]	–	+	–	–	–	–	–	10	–	–	
Area of relevé [m ²]	80	50	80	50	80	50	80	100	50	100	
Number of species in relevé	8	7	9	3	11	4	8	5	5	4	
Ch. <i>Nymphaetum candidae</i>											
<i>Nymphaea candida</i>	4	4	3	4	3	3	3	4	3	4	V
Ch. <i>Nymphaeion</i>											
<i>Myriophyllum verticillatum</i>	1	.	.	.	+	1	1	+	2	1	IV
<i>Nuphar lutea</i>	+	.	+	+	+	II
<i>Polygonum amphibium</i> fo. <i>natans</i>	+	.	+	.	+	.	1	.	.	.	II
<i>Potamogeton natans</i>	.	+	.	.	+	.	+	+	.	.	II
<i>Trapa natans</i>	+	.	+	.	1	II
Ch. <i>Potametalia, Potametea</i>											
<i>Ceratophyllum demersum</i>	+	1	+	.	+	.	+	.	.	.	III
<i>Elodea canadensis</i>	.	.	+	.	+	.	+	.	.	.	II
<i>Potamogeton pectinatus</i>	.	.	.	+	.	.	+	.	+	.	II
Ch. <i>Lemnetea minoris</i>											
<i>Lemna minor</i>	.	+	+	.	+	1	+	.	1	+	IV
<i>Spirodela polyrhiza</i>	+	+	+	.	+	+	III
<i>Salvinia natans</i>	+	.	+	.	+	.	.	.	+	.	II
<i>Utricularia vulgaris</i>	.	+	.	.	.	+	I
Accompanying species											
<i>Sphagnum fallax</i>	d	.	+	2	.	.	I
<i>Sparganium minimum</i>	.	+	+	.	.	I

Explanation: P – Przysiecz; S – Skorogoszcz; SK – Stare Kolnie; SN – Staw Nowokuźnicki Nature Reserve; So – Sowin; St – Stobrawa; SW – Stary Węgliniec; W – Wielopole; WG – Winna Góra; Z – Zielonka; Ch. – characteristic species; C – constancy; d – moss taxa.

ming in fishponds, the rush communities could disappear completely.

A threat for occurrence of *Nymphaeetum candidae* in Poland is the change of the chemical composition of water, in particular – an increase of its trophicity and hardness, which results in invasion of more expansive communities, especially of *Potametum natantis*. Conservation measures should comprise preventing eutrophication. During the fieldwork, basic threats to plant communities of the ponds were identified. The most harmful seems to be the restoration of production area of fishponds as a consequence of the program of fish culture intensification. In its result, deteriorating use of ponds is implemented, including regular mowing of the water zone and frequent deepening of ponds. The key factor allowing *Nymphaeetum candidae* to develop is temporary drying out of the ponds. If the fish farm management caused permanent filling of the pond basin with water, the studied association would probably be endangered. Localities of this association should be a subject of special protection and need to be covered with constant monitoring. The complete phytosociological and habitat characteristic as well as the present distribution of *Nymphaeetum candidae* in Poland and in the remaining countries of Central Europe, should be a subject of further botanical investigations, as this association is probably more widespread.

ACKNOWLEDGEMENTS

I wish to thank Dr. H. Ot'ahelová, Dr. R. Hrivnak (Slovak Academy of Sciences, Bratislava), Dr. D. Remy (Osnabrück University) for cooperation, Dr. M. Wayda (Jagiellonian University) for checking identification of *Nymphaea candida* and Dr. A. Stebel (Medical University of Silesia) for determination of moss taxa.

LITERATURE CITED

- BALEVIČIENĖ J., BALEVIČIUS A. 2006. Qualitative and quantitative parameters of phytocenoses in Lithuanian lakes of different trophic state. *Ekologija* 2: 34-43.
- BENKERT D., FUKAREK F., KORSCH H. 1996. Verbreitungsatlas der Farn- und Blütenpflanzen Ostdeutschland. Gustav Fischer Verlag, Jena.
- BOIŃSKI M. 1993. Pflanzenwelt des "Struga Siedmiu Jezior" Reservats in Tucheler-Heide. *Acta Univ. Nicolai Copernici, Nauki Mat.-Przyr., Biologia* 44: 93-121. (in Polish with German summary)
- BRAUN-BLANQUET J. 1964. Pflanzensoziologie, Grundzüge der Vegetationskunde. Dritte Auflage. Springer Verlag, Wien-New York.
- ČERNOHOUS F., HUSAK S. 1986. Macrophyte vegetation of Eastern and North-eastern Bohemia. *Folia Geobot. Phytotax.* 21: 113-161.
- DIERSEN K. 1996. *Vegetation Nordeuropas*. Eugen Ulmer Verlag, Stuttgart.
- FIEK E. 1881. *Flora von Schlesien*. J. U. Kern's Verl., Breslau.
- FIJAŁKOWSKI D. 1959. Plant associations of lakes situated between Łęczna and Włodawa and of peat-bogs adjacent to these lakes. *Ann. UMCS, Sec. C* 14 (3): 131-206. (in Polish with English and Russian summary)
- FRAHM J.-P., FREY W. 1992. *Moosflora*. Ed. 3. Eugen Ulmer, Stuttgart.
- HEJNÝ S. 1948. Vegetacni pomery protivinských a vodnanských rybníku. (unpublished work), Univ. Praha, Diss. (in Czech)
- HEJNÝ S. 1990. Changes in the macrophyte vegetation of South Bohemian fishponds after 35 years. *Folia Geobot. Phytotax.* 25: 245-255.
- HEJNÝ S., HUSÁK Š. 1978. Higher plant communities. In: Dykijová D., Květ J. (eds). *Pond littoral ecosystems*. Springer-Verlag, Berlin-Heidelberg-New York, pp. 23-64, 93-95.
- HEJNÝ S., SLAVIK B. 1997. *Květena České Republiky*. Vol. 1. Academia, Praha. (in Czech)
- HEREŽŇIAK J. 1972. Groupements végétaux de la vallée de la Widawka. *Monogr. Bot.* 35: 3-160. (in Polish with French summary)
- HULTEN E., FRIES M. 1986. *Atlas of North European vascular plants. North of the tropic of cancer*. Vol. 1. Koeltz Scientific Books, Koenigstein.
- JECKEL G. 1981. Die Vegetation des Naturschutzgebietes "Breites Moor" (Kreis Celle, Nordwestdeutschland). *Tuexenia* 1: 185-209.
- KŁOSOWSKI S. 2001. *Nymphaea candida* C. Presl. In: Kaźmierczakowa R., Zarzycki K. (eds). *Polish Red Data Book of Plants. Pteridophytes and flowering plants*. W. Szafer Institute of Botany, Instyute of Nature Conservation, Polish Academy of Sciences, Cracow, pp. 113-116. (in Polish with English summary)
- KRZYWAŃSKI D. 1974. The plant communities of old riverbeds of middle part of the Warta river in Central Poland. *Monogr. Bot.* 43: 1-80. (in Polish with English summary)
- KUBÁT K., HROUDA L., CHRTEK J. JUN., KAPLAN Z., KIRSCHNER J., ŠTĚPÁNEK J. (eds) 2002. *Klíč ke květeně České republiky*. Academia, Praha. (in Czech)
- MACICKA-PAWLIK T., WILCZYŃSKA W. 1996. Plant communities of old river-beds in the valley of the Odra river middle course. *Acta Univ. Wratisl. 1735, Pr. Bot.* 64: 72-120. (in Polish with English summary)
- MAMIŃSKI M. 1986. Peatbog plant communities of Bełchatów Industrial Region. *Acta Univ. Lodz., Folia Bot.* 4: 85-137. (in Polish with English summary)
- MATUSZKIEWICZ W. 2005. *Przewodnik do oznaczania zbiorowiska roślinnych Polski*. Wydawnictwo Naukowe PWN, Warszawa. (in Polish)
- MEUSEL H., JÄGER E., WEINERT E. 1965. *Vergleichende Chorologie der Zentraleuropäischen Flora*. Gustav Fischer Verlag, Jena.
- MILJAN A. 1933. Vegetationsuntersuchungen an Naturwiesen und Seen im Osteuropäischen Moränengebiet Estlands. *Acta Comment.* 25: 1-132.
- MIREK Z., PIĘKOŚ-MIRKOWA H., ZAJĄC A., ZAJĄC M. 2002. Flowering plants and pteridophytes of Poland – a checklist. In: Mirek Z. (ed.). *Biodiversity of Poland I*, W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków, p. 442.
- MUNTENDAM J. B., POVEL G.D.E., VAN DER VELDE G. 1996. Morphometric patterns in the *Nymphaea alba-candida* complex. *Acta Bot. Neerl.* 45 (3): 279-302.
- NEUHÄUSL R., NEUHÄUSLOVA Z. 1965. Die Pflanzengesellschaften des Naturschutzgebietes "Berehynsky Rybnik" bei Doksy (Hirschberg). *Preslia* 37: 170-199.
- NEUHÄUSL R., TOMSOVIC P. 1957. Die Gattung *Nymphaea* (L.) Smith in der Tschechoslowakei. *Preslia* 29: 225-249.
- NOWAK A., NOWAK S. 2007. The problem of the occurrence of *Nymphaea candida* C. Presl in the Opole Silesia. *Opole Scientific Society Nature Journal* 40: 5-16.
- OBERDORFER E. 1994. *Pflanzensoziologische Exkursionsflora*. 7 Aufl. Verlag Eugen Ulmer, Stuttgart.
- PASSARGE H. 1957. Über Wasserpflanzen- und Kleinröhrichtgesellschaften des Oberspreewaldes. *Abh. Ber. Naturk. Mus. Görlitz* 35: 143-152.
- PASSARGE H. 1996. *Pflanzengesellschaften Nordostdeutschlands*. I. Hydro- und Therophytosa. J. Cramer, Berlin-Stuttgart.

- PODPIELKOWSKI Z. 1968. Die Vegetation der Fischteiche der Woiwodschaft Warszawa. Monogr. Bot. 27: 3-122. (in Polish with German summary)
- POTT R. 1995. Die Pflanzengesellschaften Deutschlands. 2. Aufl. Eugen Ulmer Verlag, Stuttgart.
- RENNWALD E. (ed.) 2000. Rote Liste der Pflanzengesellschaften Deutschlands mit Anmerkungen zur Gefährdung. In: Rennwald E. (ed.). Verzeichnis und Rote Liste der Pflanzengesellschaften Deutschlands. Schr.-R.f. Vegetationskunde 35: 393-592.
- SCHUBE T. 1903. Die Verbreitung der Gefäßpflanzen in Schlesien, preussischen und österreichischen Anteils. Druck von R. Nischowsky, Breslau.
- SCHUBERT R., HILBIG W., KLOTZ S. 1995. Bestimmungsbuch der Pflanzengesellschaften Mittel- und Nordostdeutschlands. G. Fischer Verlag, Jena-Stuttgart.
- SPAŁEK K. 2004. The state, threat and conservation of the water, rush and alluvium vegetation. In: A. Nowak, K. Spałek (eds). Conservation of vegetation of the Opole Silesia. Uniwersytet Opolski, Opole, pp. 227-234. (in Polish with English abstract)
- SPAŁEK K. 2005. Scirpetum radicans Hejný in Hejný et Husák 1978 in Poland. Thaiszia 15: 43-51.
- SPAŁEK K. 2006a. Threatened plant communities as an indicator of fishponds value: an example from Silesia (SW Poland). In: Gafta D., Akeroyd J. (eds). Nature Conservation. Concepts and Practice. Springer Verlag, Berlin-Heidelberg, pp. 195-198.
- SPAŁEK K. 2006b. Utricularietum australis Th. Müller et Görs 1960 in Poland. Acta Soc. Bot. Pol. 75 (3): 253-256.
- SPAŁEK K. 2007. Nymphaetum albae Vollmar 1947 em. Oberd. in Oberd. et al. 1967, a plant association new to Poland. Acta Soc. Bot. Pol. 76 (1): 81-84.
- SUGIER P., POPIOŁEK Z. 1995. The aquatic and coastal vegetation of Polesie National Park against the background of habitat conditions. Lake Karaśne. Ann. UMCS, Sec. C 50: 55-69. (in Polish with English summary)
- SUGIER P., POPIOŁEK Z. 1998. The aquatic and coastal vegetation of lake Moszne in Polesie National Park. Ann. UMCS, Sec. C 53: 185-200. (in Polish with English summary)
- SZUMIEC M.A. 1995. The role of carp pond ecosystem management in wetland conservation. Acta Hydrobiol. 37, Suppl. 1: 13-20.
- TOMASZEWICZ H. 1977. Proposal of new syntaxonomic classification of Myriophyllo-Nupharetum W. Koch 1926 phytocoenoses and their distribution in Poland. Acta Soc. Bot. Pol. 46 (3): 423-436.
- TOMASZEWICZ H. 1979. Roślinność wodna i szuwarowa Polski (Klasy: Lemnetaea, Charetea, Potamogetonetea, Phragmitetea) wg stanu zbadania na rok 1975. Rozpr. Uniw. Warsz., Warszawa. (in Polish)
- TOMASZEWICZ H., KŁOSOWSKI S. 1993. Standortsverhältnisse der Gesellschaften mit Dominanz einzelner Nymphaeaceen in Nordost-Polen. Tuexenia 13: 75-90.
- TUTIN T.G., BURGESS N.A., CHATER A.O., EDMONDSON J.R., HEYWOOD V.E., MOORE D.M., VALENTINE D.H., WALTERS S.M., WEBB D.A. (ed.) 2002. Flora Europaea. Vol. 1. Cambridge University Press, Cambridge.
- WAYDA M. 2000. The distribution of *Nymphaea candida* C. Presl (Nymphaeaceae) in Poland. Acta Soc. Bot. Pol. 69 (1): 75-78.
- WEBER-OLDECOP D.W. 1975. Die glänzendweiße Seerose (*Nymphaea candida* Presl.) in der Lüneburger Heide. Gött. Flor. Rundbr. 9 (3): 86-87.
- WOJTASZEK M. 1989. Vegetation of Old River-Beds on the Right Bank of Warta Valley in the Region of Rogalin. Bad. Fizjogr. Pol. Zach., Ser. B 39: 105-117. (in Polish with English summary)
- ZABAŃSKI J., MATUŁA J. 1973. *Nymphaea candida* in the district of Zgorzelec on the Lower Silesia. Chrońmy Przyr. Ojcz. 29 (1): 76-80. (in Polish)
- ZAJĄC A., ZAJĄC M. (ed.) 2001. Distribution Atlas of Vascular Plants in Poland. Edited by Laboratory of Computer Chorology, Institute of Botany, Jagiellonian University, Cracow.